#### HK j efacean research project 香港鯨豚研究計劃

#### HK CETACEAN RESEARCH PROJECT

### 香港鯨豚研究計劃

#### CONTRACT NO. HY/2012/07

# Hong Kong-Zhuhai-Macao Bridge Tuen Mun – Chek Lap Kok Link (Southern Connection Viaduct Section) Dolphin Quarterly Monitoring

16<sup>th</sup> Quarterly Progress Report (September-November 2017) submitted to Gammon Construction Limited

Submitted by Samuel K.Y. Hung, Ph.D., Hong Kong Cetacean Research Project

13 March 2018

#### 1. Introduction

- 1.1. The Tuen Mun-Chek Lap Kok Link (TM-CLKL) comprises a 1.6 km long dual 2-lane viaduct section between the Hong Kong Boundary Crossing Facilities (HKBCF) and the North Lantau Highway and associated roads at Tai Ho. Gammon Construction Limited (hereinafter called the "Contractor") was awarded as the main contractor of "Contract No. HY/2012/07 Hong Kong-Zhuhai-Macao Bridge Tuen Mun-Chek Lap Kok Link Southern Connection Viaduct Section".
- 1.2. According to the updated Environmental Monitoring and Audit (EM&A) Manual (for TM-CLKL), monthly line-transect vessel surveys for Chinese White Dolphin should be conducted to cover the Northwest (NWL) and Northeast Lantau (NEL) survey areas as in AFCD annual marine mammal monitoring programme. However, as such surveys have been undertaken by the HKLR03 and HKBCF projects in the same areas (i.e. NWL and NEL), a combined monitoring approach is recommended by the Highways Department, that the TM-CLKL EM&A project can utilize the monitoring data collected by HKLR03 or HKBCF project to avoid any redundancy in monitoring effort. Such exemption for the dolphin monitoring will end upon the completion of the dolphin monitoring carried out by HKLR03 contract as well as the TM-CLKL Northern Connection Sub-Sea Tunnel Section (HY/2012/08).
- 1.3. In November 2013, the Director of Hong Kong Cetacean Research Project (HKCRP), Dr. Samuel Hung, has been appointed by Gammon Construction Limited as the dolphin specialist for the TM-CLKL Southern Viaduct Section EM&A project. He is responsible for the dolphin monitoring study, including the data collection on Chinese White Dolphins during the construction phase (i.e. impact period) of the TM-CLKL project in Northwest Lantau (NWL) and Northeast Lantau (NEL) survey areas.
- 1.4. During the construction period of HKLR, the dolphin specialist would be in charge of



## 香港鯨豚研究計劃

reviewing and collating information collected by the HKLR03 dolphin monitoring programme to examine any potential impacts of TM-CLKL construction works on the dolphins.

- 1.5. From the monitoring results, any changes in dolphin occurrence within the study area will be examined for possible causes, and appropriate actions and additional mitigation measures will be recommended as necessary.
- 1.6. This report is the 16<sup>th</sup> quarterly progress report under the TM-CLKL construction phase dolphin monitoring programme submitted to the Gammon Construction Limited, summarizing the results of the surveys findings during the period of September to November 2017, utilizing the survey data collected by HKLR03 impact phase monitoring project.

#### 2. Monitoring Methodology

- 2.1. Vessel-based Line-transect Survey
- 2.1.1. According to the requirement of the updated EM&A manual, dolphin monitoring programme should cover all transect lines in NEL and NWL survey areas (see Figure 1) twice per month throughout the entire construction period. The co-ordinates of all transect lines are shown in Table 1.

Table 1 Co-ordinates of transect lines conducted by HKLR03 project

	Line No.	Easting	Northing	Line No.		Easting	Northing
1	Start Point	804671	815456	13	Start Point	816506	819480
1	End Point	804671	831404	13	End Point	816506	824859
2	Start Point	805476	820800	14	Start Point	817537	820220
2	End Point	805476	826654	14	End Point	817537	824613
3	Start Point	806464	821150	15	Start Point	818568	820735
3	End Point	806464	822911	15	End Point	818568	824433
4	Start Point	807518	821500	16	Start Point	819532	821420
4	End Point	807518	829230	16	End Point	819532	824209
5	Start Point	808504	821850	17	Start Point	820451	822125
5	End Point	808504	828602	17	End Point	820451	823671
6	Start Point	809490	822150	18	Start Point	821504	822371
6	End Point	809490	825352	18	End Point	821504	823761
7	Start Point	810499	822000	19	Start Point	822513	823268
7	End Point	810499	824613	19	End Point	822513	824321
8	Start Point	811508	821123	20	Start Point	823477	823402



## 香港鯨豚研究計劃

8	End Point	811508	824254	20	End Point	823477	824613
9	Start Point	812516	821303	21	Start Point	805476	827081
9	End Point	812516	824254	21	End Point	805476	830562
10	Start Point	813525	821176	22	Start Point	806464	824033
10	End Point	813525	824657	22	End Point	806464	829598
11	Start Point	814556	818853	23	Start Point	814559	821739
11	End Point	814556	820992	23	End Point	814559	824768
12	Start Point	815542	818807	24	Start Point	805476	815900
12	End Point	815542	824882	24	End Point	805476	819100

- 2.1.2. The HKLR03 survey team used standard line-transect methods (Buckland et al. 2001) to conduct the systematic vessel surveys, and followed the same technique of data collection that has been adopted over the last 20 years of marine mammal monitoring surveys in Hong Kong developed by HKCRP (see Hung 2017). For each monitoring vessel survey, a 15-m inboard vessel with an open upper deck (about 4.5 m above water surface) was used to make observations from the flying bridge area.
- 2.1.3. Two experienced observers (a data recorder and a primary observer) made up the on-effort survey team, and the survey vessel transited different transect lines at a constant speed of 13-15 km per hour. The data recorder searched with unaided eyes and filled out the datasheets, while the primary observer searched for dolphins and porpoises continuously through 7 x 50 *Fujinon* marine binoculars. Both observers searched the sea ahead of the vessel, between 270° and 90° (in relation to the bow, which is defined as 0°). One to two additional experienced observers were available on the boat to work in shift (i.e. rotate every 30 minutes) in order to minimize fatigue of the survey team members. All observers were experienced in small cetacean survey techniques and identifying local cetacean species.
- 2.1.4. During on-effort survey periods, the survey team recorded effort data including time, positions (latitude and longitude), weather conditions (Beaufort sea state and visibility), and distance traveled in each series (a continuous period of search effort) with the assistance of a handheld GPS (*Garmin eTrex Legend*).
- 2.1.5. Data including time, position and vessel speed were also automatically and continuously logged by handheld GPS throughout the entire survey for subsequent review.
- 2.1.6. When dolphins were sighted, the survey team would end the survey effort, and immediately record the initial sighting distance and angle of the dolphin group from the survey vessel, as well as the sighting time and position. Then the research vessel was diverted from its course to approach the animals for species identification, group size estimation, assessment of group composition, and behavioural observations. The perpendicular distance (PSD) of the dolphin group to the transect line was later calculated from the initial sighting distance and angle.



2.1.7. Survey effort being conducted along the parallel transect lines that were perpendicular to the coastlines (as indicated in Figure 1) was labeled as "primary" survey effort, while the survey effort conducted along the connecting lines between parallel lines was labeled as "secondary" survey effort. According to HKCRP long-term dolphin monitoring data, encounter rates of Chinese white dolphins deduced from effort and sighting data collected along primary and secondary lines were similar in NEL and NWL survey areas. Therefore, both primary and secondary survey effort were presented as on-effort survey effort in this report.

#### 2.2. Photo-identification Work

- 2.2.1. When a group of Chinese White Dolphins were sighted during the line-transect survey, the HKLR03 survey team would end effort and approach the group slowly from the side and behind to take photographs of them. Every attempt was made to photograph every dolphin in the group, and even photograph both sides of the dolphins, since the colouration and markings on both sides may not be symmetrical.
- 2.2.2. A professional digital camera (*Canon* EOS 7D model), equipped with long telephoto lenses (100-400 mm zoom), were available on board for researchers to take sharp, close-up photographs of dolphins as they surfaced. The images were shot at the highest available resolution and stored on Compact Flash memory cards for downloading onto a computer.
- 2.2.3. All digital images taken in the field were first examined, and those containing potentially identifiable individuals were sorted out. These photographs would then be examined in greater detail, and were carefully compared to the existing Chinese White Dolphin photo-identification catalogue maintained by HKCRP since 1995.
- 2.2.4. Chinese White Dolphins can be identified by their natural markings, such as nicks, cuts, scars and deformities on their dorsal fin and body, and their unique spotting patterns were also used as secondary identifying features (Jefferson 2000).
- 2.2.5. All photographs of each individual were then compiled and arranged in chronological order, with data including the date and location first identified (initial sighting), re-sightings, associated dolphins, distinctive features, and age classes entered into a computer database.

#### 2.3. Data Analysis

- 2.3.1. Distribution Analysis The line-transect survey data was integrated with the Geographic Information System (GIS) in order to visualize and interpret different spatial and temporal patterns of dolphin distribution using sighting positions. Location data of dolphin groups were plotted on map layers of Hong Kong using a desktop GIS (ArcView<sup>©</sup> 3.1) to examine their distribution patterns in details. The dataset was also stratified into different subsets to examine distribution patterns of dolphin groups with different categories of group sizes, young calves and activities.
- 2.3.2. Encounter rate analysis Encounter rates of Chinese white dolphins (number of on-effort



### 香港鯨豚研究計劃

sightings per 100 km of survey effort, and total number of dolphins sighted on-effort per 100 km of survey effort) were calculated in NEL and NWL survey areas in relation to the amount of survey effort conducted during each month of monitoring survey. Only data collect under Beaufort 3 or below condition would be used for the encounter rate analyses. Dolphin encounter rates were calculated in two ways for comparisons with the HZMB baseline monitoring results as well as to AFCD long-term marine mammal monitoring results.

Firstly, for the comparison with the HZMB baseline monitoring results, the encounter rates were calculated using primary survey effort alone. The average encounter rate of sightings (STG) and average encounter rate of dolphins (ANI) were deduced based on the encounter rates from six events during the present quarter (i.e. six sets of line-transect surveys in North Lantau), which was also compared with the one deduced from the six events during the baseline period (i.e. six sets of line-transect surveys in North Lantau).

Secondly, the encounter rates were calculated using both primary and secondary survey effort collected under Beaufort 3 or below condition as in AFCD long-term monitoring study. The encounter rate of sightings and dolphins were deduced by dividing the total number of on-effort sightings (STG) and total number of dolphins (ANI) by the amount of survey effort for the present quarterly period.

2.3.3. Quantitative grid analysis on habitat use – To conduct quantitative grid analysis of habitat use, positions of on-effort sightings of Chinese White Dolphins collected during the quarterly impact phase monitoring period were plotted onto 1-km² grids among NWL and NEL survey areas on GIS. Sighting densities (number of on-effort sightings per km²) and dolphin densities (total number of dolphins from on-effort sightings per km²) were then calculated for each 1 km by 1 km grid with the aid of GIS. Sighting density grids and dolphin density grids were then further normalized with the amount of survey effort conducted within each grid. The total amount of survey effort spent on each grid was calculated by examining the survey coverage on each line-transect survey to determine how many times the grid was surveyed during the study period. For example, when the survey boat traversed through a specific grid 50 times, 50 units of survey effort were counted for that grid. With the amount of survey effort calculated for each grid, the sighting density and dolphin density of each grid were then normalized (i.e. divided by the unit of survey effort).

The newly-derived unit for sighting density was termed SPSE, representing the number of on-effort sightings per 100 units of survey effort. In addition, the derived unit for actual dolphin density was termed DPSE, representing the number of dolphins per 100 units of survey effort. Among the 1-km² grids that were partially covered by land, the percentage of sea area was calculated using GIS tools, and their SPSE and DPSE values were adjusted accordingly. The following formulae were used to estimate SPSE and DPSE in each 1-km² grid within the study area:

SPSE = ((S / E) x 100) / SA% DPSE = ((D / E) x 100) / SA%

where S = total number of on-effort sightings



## 香港鯨豚研究計劃

D = total number of dolphins from on-effort sightings E = total number of units of survey effort SA% = percentage of sea area

- 2.3.4. Behavioural analysis When dolphins were sighted during vessel surveys, their behaviour was observed. Different activities were categorized (i.e. feeding, socializing, traveling, and milling/resting) and recorded on sighting datasheets. This data was then input into a separate database with sighting information, which can be used to determine the distribution of behavioural data with a desktop GIS. Distribution of sightings of dolphins engaged in different activities and behaviours would then be plotted on GIS and carefully examined to identify important areas for different activities of the dolphins.
- 2.3.5. Ranging pattern analysis Location data of individual dolphins that occurred during the 3-month impact phase monitoring period were obtained from the dolphin sighting database and photo-identification catalogue. To deduce home ranges for individual dolphins using the fixed kernel methods, the program Animal Movement Analyst Extension, was loaded as an extension with ArcView® 3.1 along with another extension Spatial Analyst 2.0. Using the fixed kernel method, the program calculated kernel density estimates based on all sighting positions, and provided an active interface to display kernel density plots. The kernel estimator then calculated and displayed the overall ranging area at 95% UD level.

#### 3. Monitoring Results

- 3.1. Summary of survey effort and dolphin sightings
- 3.1.1. During the period of September to November 2017, six sets of systematic line-transect vessel surveys were conducted under the HKLR03 monitoring works to cover all transect lines in NWL and NEL survey areas twice per month.
- 3.1.2. From these HKLR03 surveys, a total of 802.12 km of survey effort was collected, with 96.0% of the total survey effort being conducted under favourable weather conditions (i.e. Beaufort Sea State 3 or below with good visibility). Among the two areas, 297.00 km and 505.12 km of survey effort were conducted in NEL and NWL survey areas respectively.
- 3.1.3. The total survey effort conducted on primary lines was 578.16 km, while the effort on secondary lines was 223.96 km. Survey effort conducted on both primary and secondary lines were considered as on-effort survey data. A summary table of the survey effort is shown in Appendix I.
- 3.1.4. During the six sets of HKLR03 monitoring surveys from September to November 2017, 13 groups of 50 Chinese White Dolphins were sighted. All dolphin sightings were made during on-effort search in this quarter, and 12 of the 13 on-effort dolphin sightings were made on primary lines. A summary table of dolphin sightings is shown in Appendix II.



- 3.1.5. In this quarterly period, all dolphin groups were sighted in NWL, and no dolphin was sighted at all in NEL. In fact, since August 2014, only two sightings of two lone dolphins were made respectively in NEL during HKLR03 monitoring surveys.
- 3.2. Distribution
- 3.2.1. Distribution of dolphin sightings made during the HKLR03 monitoring surveys from September to November 2017 is shown in Figure 1. Almost all sightings were made at the northwest portion of the North Lantau region, mainly to the east of Lung Kwu Chau and at the mouth of Deep Bay near Black Point (Figure 1). One dolphin group was also sighted at the southwestern end of NWL survey area, or near the HKLR09 alignment. As consistently recorded in the previous monitoring quarters, the dolphins were completely absent from the central and eastern portions of North Lantau waters (Figure 1).
- 3.2.2. All dolphin sightings were located far away from the alignments of TM-CLKL as well as the HKBCF and HKLR03 reclamation sites (Figure 1). However, one dolphin group was sighted near the alignment of HKLR09 as mentioned above.
- 3.2.3. Sighting distribution of dolphins during the present impact phase monitoring period (September to November 2017) was drastically different from the one during the baseline monitoring period (Figure 1). In the present quarter, dolphins have disappeared from the NEL region, which was in stark contrast to their frequent occurrence around the Brothers Islands, near Shum Shui Kok and in the vicinity of HKBCF reclamation site during the baseline period (Figure 1). The nearly complete abandonment of NEL region by the dolphins has been consistently recorded in the past 18 quarters of HKLR03 monitoring, which has resulted in zero to extremely low dolphin encounter rates in this area.
- 3.2.4. In NWL survey area, dolphin occurrence was also significantly different between the baseline and impact phase periods. During the present impact monitoring period, dolphins were infrequently sighted here, and mainly at the northwestern end of the area, which was in stark contrast with their frequent occurrences throughout the area during the baseline period (Figure 1).
- 3.2.5. Another comparison in dolphin distribution was made between the five quarterly periods of autumn months in 2013-17 (Figure 2). Among the five autumn periods, dolphins were still sighted regularly in NWL waters in 2013 and 2014, but their usage there was progressively reduced in the three subsequent autumn periods, with the only occurrences mostly concentrated at the northwestern portion of the survey area (Figure 2).
- 3.3. Encounter rate
- 3.3.1. During the present quarterly period, the encounter rates of Chinese White Dolphins deduced from the survey effort and on-effort sighting data from the primary transect lines under favourable conditions (Beaufort 3 or below) for each set of the HKLR03 surveys in NEL and NWL are shown in Table 2. The average encounter rates deduced from the six sets of HKLR03 surveys were also compared with the ones deduced from the baseline monitoring period (September November 2011) (Table 3).



## 香港鯨豚研究計劃

Table 2. Dolphin encounter rates (sightings per 100 km of survey effort) during September-November 2017

SURVEY AREA	DOLPHIN MONITORING DATES	Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of survey effort)  Primary Lines Only	Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort)  Primary Lines Only		
	Set 1 (15 & 18 Sep 2017)	0.00	0.00		
	Set 2 (22 & 29 Sep 2017)	0.00	0.00		
Northeast	Set 3 (4 & 9 Oct 2017)	0.00	0.00		
Lantau	Set 4 (18 & 26 Oct 2017)	0.00	0.00		
	Set 5 (1 & 8 Nov 2017)	0.00	0.00		
	Set 6 (17 & 24 Nov 2017)	0.00	0.00		
	Set 1 (15 & 18 Sep 2017)	0.00	0.00		
	Set 2 (22 & 29 Sep 2017)	3.63	16.34		
Northwest	Set 3 (4 & 9 Oct 2017)	1.86	9.30		
Lantau	Set 4 (18 & 26 Oct 2017)	4.89	4.89		
	Set 5 (1 & 8 Nov 2017)	4.99	26.60		
	Set 6 (17 & 24 Nov 2017)	3.33	5.00		

Table 3. Comparison of average dolphin encounter rates from impact monitoring period (September - November 2017) and baseline monitoring period (September – November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions; ± denotes the standard deviation of the average encounter rates)

	Encounter I (no. of on-effort dolph km of surve	in sightings per 100	Encounter rate (ANI) (no. of dolphins from all on-effort sightings per 100 km of survey effort)		
	September – November 2017	September – November 2011	September – November 2017	September – November 2011	
Northeast Lantau	0.0	6.00 ± 5.05	0.0	22.19 ± 26.81	
Northwest Lantau	3.12 ± 1.91	9.85 ± 5.85	10.35 ± 9.66	44.66 ± 29.85	

- 3.3.2. To facilitate the comparison with the AFCD long-term monitoring results, the encounter rates were also calculated for the present quarter using both primary and secondary survey effort. The encounter rates of sightings (STG) and dolphins (ANI) in NWL were 2.5 sightings and 9.9 dolphins per 100 km of survey effort respectively, while the encounter rates of sightings (STG) and dolphins (ANI) in NEL were both nil for this quarter.
- 3.3.3. In NEL, the average dolphin encounter rates (both STG and ANI) in the present three-month impact monitoring period were both zero with no on-effort sighting being made, and such extremely low occurrence of dolphins in NEL have been consistently

#### HK J efacean research project 香港鯨豚研究計劃

#### HK CETACEAN RESEARCH PROJECT

### 香港鯨豚研究計劃

recorded in the past 18 quarters of HKLR03 monitoring (Table 4). This is a serious concern as the dolphin occurrence in NEL in the past few years (0.0-1.0 for ER(STG) and 0.0-3.9 for ER(ANI)) have remained exceptionally low when compared to the baseline period (Table 4). Dolphins have been virtually absent from NEL waters since January 2014, with only three groups of six dolphins sighted there since then despite consistent and intensive survey effort being conducted in this survey area.

Table 4. Comparison of average dolphin encounter rates in Northeast Lantau survey area from all quarters of HKLR03 impact monitoring period and baseline monitoring period (September-November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions; the encounter rates in **autumn** months were highlighted in **blue**; ± denotes the standard deviation of the average encounter rates)

	Encounter rate (STG) (no. of on-effort dolphin sightings per 100 km of	(no. of dolphins from all on-effort sightings per 100
September-November 2011 (Baseline)	survey effort) 6.00 ± 5.05	km of survey effort)  22.19 ± 26.81
December 2012-February 2013 (Impact)	3.14 ± 3.21	6.33 ± 8.64
March-May 2013 (Impact)	0.42 ± 1.03	0.42 ± 1.03
June-August 2013 (Impact)	0.88 ± 1.36	3.91 ± 8.36
September-November 2013 (Impact)	1.01 ± 1.59	3.77 ± 6.49
December 2013-February 2014 (Impact)	0.45 ± 1.10	1.34 ± 3.29
March-May 2014 (Impact)	0.00	0.00
June-August 2014 (Impact)	0.42 ± 1.04	1.69 ± 4.15
September-November 2014 (Impact)	0.00	0.00
December 2014-February 2015 (Impact)	0.00	0.00
March-May 2015 (Impact)	0.00	0.00
June-August 2015 (Impact)	0.44 ± 1.08	0.44 ± 1.08
September-November 2015 (Impact)	0.00	0.00
December 2015-February 2016 (Impact)	0.00	0.00
March-May 2016 (Impact)	0.00	0.00
June-August 2016 (Impact)	0.00	0.00
September-November 2016 (Impact)	0.00	0.00
December 2016-February 2017 (Impact)	0.00	0.00
March-May 2017 (Impact)	0.00	0.00
June-August 2017 (Impact)	0.00	0.00
September-November 2017 (Impact)	0.00	0.00

3.3.4. On the other hand, the average dolphin encounter rates (STG and ANI) in NWL during



### 香港鯨豚研究計劃

the present impact phase monitoring period (reductions of 68.3% and 76.8% respectively) were only small fractions of the ones recorded during the three-month baseline period, indicating a dramatic decline in dolphin usage of this survey area as well during the present impact phase period (Table 5).

Table 5. Comparison of average dolphin encounter rates in Northwest Lantau survey area from all quarters of HKLR03 impact monitoring period and baseline monitoring period (September-November 2011) (Note: encounter rates deduced from the baseline monitoring period have been recalculated based only on survey effort and on-effort sighting data made along the primary transect lines under favourable conditions; the encounter rates in **autumn** months were highlighted in **blue**; ± denotes the standard deviation of the average encounter rates)

	Encounter rate (STG) (no. of on-effort dolphin	Encounter rate (ANI) (no. of dolphins from all
	sightings per 100 km of	on-effort sightings per 100
	survey effort)	km of survey effort)
September-November 2011 (Baseline)	9.85 ± 5.85	44.66 ± 29.85
December 2012-February 2013 (Impact)	8.36 ± 5.03	35.90 ± 23.10
March-May 2013 (Impact)	7.75 ± 3.96	24.23 ± 18.05
June-August 2013 (Impact)	6.56 ± 3.68	27.00 ± 18.71
September-November 2013 (Impact)	8.04 ± 1.10	32.48 ± 26.51
December 2013-February 2014 (Impact)	8.21 ± 2.21	32.58 ± 11.21
March-May 2014 (Impact)	6.51 ± 3.34	19.14 ± 7.19
June-August 2014 (Impact)	4.74 ± 3.84	17.52 ± 15.12
September-November 2014 (Impact)	5.10 ± 4.40	20.52 ± 15.10
December 2014-February 2015 (Impact)	2.91 ± 2.69	11.27 ± 15.19
March-May 2015 (Impact)	0.47 ± 0.73	2.36 ± 4.07
June-August 2015 (Impact)	2.53 ± 3.20	9.21 ± 11.57
September-November 2015 (Impact)	3.94 ± 1.57	21.05 ± 17.19
December 2015-February 2016 (Impact)	2.64 ± 1.52	10.98 ± 3.81
March-May 2016 (Impact)	0.98 ± 1.10	4.78 ± 6.85
June-August 2016 (Impact)	1.72 ± 2.17	7.48 ± 10.98
September-November 2016 (Impact)	2.86 ± 1.98	10.89 ± 10.98
December 2016-February 2017 (Impact)	3.80 ± 3.79	14.52 ± 17.21
March-May 2017 (Impact)	0.93 ± 1.03	5.25 ± 9.53
June-August 2017 (Impact)	2.20 ± 2.88	6.58 ± 8.12
September-November 2017 (Impact)	3.12 ± 1.91	10.35 ± 9.66

3.3.5. Dolphin encounter rates in NWL during autumn 2017 was similar to the previous autumn period in 2016, but was much lower than the ones in the autumn periods of 2013, 2014 and 2015 (Table 5). Such temporal trend should be closely monitored in the upcoming



monitoring quarters whether the dolphin occurrence would continue to increase as the construction activities of HZMB works have been mostly completed in coming months.

- 3.3.6 As discussed in Hung (2017), the dramatic decline in dolphin usage of NEL waters in the past few years (including the declines in abundance, encounter rate and habitat use in NEL, as well as shifts of individual core areas and ranges away from NEL waters) was possibly related to the HZMB construction works that were commenced since 2012. Apparently such noticeable decline has already extended to NWL waters progressively in the past few years with no sign of recovery, even though the HZMB-related construction activities have well past the peak.
- 3.3.7. A two-way ANOVA with repeated measures and unequal sample size was conducted to examine whether there were any significant differences in the average encounter rates between the baseline and impact monitoring periods. The two variables that were examined included the two periods (baseline and impact phases) and two locations (NEL and NWL).
- 3.3.8. For the comparison between the baseline period and the present quarter (20<sup>th</sup> quarter of the HKLR03 impact phase being assessed), the p-values for the differences in average dolphin encounter rates of STG and ANI were 0.0057 and 0.0278 respectively. If the alpha value is set at 0.05, significant differences were detected between the baseline and present quarters in both the average dolphin encounter rates of STG and ANI.
- 3.3.9. For the comparison between the baseline period and the cumulative quarters in impact phase (i.e. the first 20 quarters of the HKLR03 impact phase being assessed), the p-values for the differences in average dolphin encounter rates of STG and ANI were 0.000000 and 0.000000 respectively. Even if the alpha value is set at 0.00001, significant differences were still detected in both the average dolphin encounter rates of STG and ANI (i.e. between the two periods and the locations).
- 3.3.10. As indicated in both dolphin distribution patterns and encounter rates, dolphin usage has been significantly reduced in both NEL and NWL survey areas during the present quarterly period, and such low occurrence of dolphins has also been consistently documented in previous quarters of the past few years.
- 3.3.11. The dramatic decline in dolphin usage of North Lantau region raises serious concern, as the timing of the decline in dolphin usage in North Lantau waters coincided well with the construction schedule of the HZMB-related projects (Hung 2017). Apparently there was no sign of recovery of dolphin usage even though almost all marine works associated with the HZMB construction have been completed.
- 3.4. Group size
- 3.4.1. Group size of Chinese White Dolphins ranged from one to 12 individuals per group in North Lantau region during September to November 2017. The average dolphin group sizes from these three months were compared with the ones deduced from the baseline period in September to November 2011, as shown in Table 6.



## 香港鯨豚研究計劃

3.4.2. The average dolphin group size in NWL waters during September to November 2017 was only slightly higher than the one recorded during the three-month baseline period, but it should also be noted that the sample size of 13 dolphin groups in the present quarter was very small when compared to the 66 groups sighted during the baseline period (Table 6).

Table 6. Comparison of average dolphin group sizes from impact monitoring period (September – November 2017) and baseline monitoring period (September – November 2011) (Note: ± denotes the standard deviation of the average group size)

	Average Dolphin Group Size							
	September – November 2017	September – November 2011						
Overall	3.85 ± 3.39 (n = 13)	3.72 ± 3.13 (n = 66)						
Northeast Lantau		3.18 ± 2.16 (n = 17)						
Northwest Lantau	$3.85 \pm 3.39 (n = 13)$	3.92 ± 3.40 (n = 49)						

- 3.4.3. Notably, 8 of these 14 dolphin groups were composed of 1-3 individuals only, while there were four medium-sized groups with 5-8 dolphins per group, and one large group of 12 dolphins (Appendix II).
- 3.4.4. Distribution of the larger dolphin groups with five individuals or more per group during the present quarter is shown in Figure 3, with comparison to the one in baseline period. The medium-sized group with 5-8 dolphins were scattered at the northwestern portion of the NWL survey area with no particular concentration, while the one large group of 12 dolphins was sighted at the mouth of Deep Bay (Figure 3). Such distribution pattern was very different from the baseline period, when the larger dolphin groups were frequently sighted and evenly distributed in NWL waters, and a few were also sighted in NEL waters (Figure 3).

#### 3.5. Habitat use

- 3.5.1. From September to November 2017, four of the five grids with moderately high to high dolphin densities were located to the north of Lung Kwu Chau, while one grid to the east of Sha Chau also recorded moderately high dolphin density (Figures 4a and 4b). All grids near HKLR03/HKBCF reclamation sites as well as TMCLKL alignment did not record any presence of dolphins at all during on-effort search in the present quarterly period (Figures 4a and 4b).
- 3.5.2. However, it should be emphasized that the amount of survey effort collected in each grid during the three-month period was fairly low (6-12 units of survey effort for most grids), and therefore the habitat use pattern derived from the three-month dataset should be treated with caution. A more complete picture of dolphin habitat use pattern should be examined when more survey effort for each grid will be collected throughout the impact phase monitoring programme.
- 3.5.3. When compared with the habitat use patterns during the baseline period, dolphin usage in NEL and NWL has drastically diminished in both areas during the present impact monitoring period (Figure 5). During the baseline period, many grids between Siu Mo



## 香港鯨豚研究計劃

To and Shum Shui Kok in NEL recorded moderately high to high dolphin densities, which was in stark contrast to the complete absence of dolphins there during the present impact phase period (Figure 5).

- 3.5.4. The density patterns were also very different in NWL between the baseline and impact phase monitoring periods, with high dolphin usage throughout the area, especially around Sha Chau, near Black Point, to the west of the airport, as well as between Pillar Point and airport platform during the baseline period. In contrast, only several grids with moderately high to high dolphin densities were located near Lung Kwu Chau and Sha Chau during the present impact phase period (Figure 5).
- 3.6. Mother-calf pairs
- 3.6.1. During the present quarterly period, no young calf was sighted at all among the 13 groups of dolphins.
- 3.7. Activities and associations with fishing boats
- 3.7.1. One of the thirteen dolphin groups were engaged in feeding activity, while another two groups were engaged in socializing activity. However, none of them was engaged in traveling or milling/resting activity during the three-month study period.
- 3.7.2. The percentages of sightings associated with feeding activities (7.7%) was lower than the one recorded during the baseline period (11.6%), while the one for socializing activities (15.4%) was much higher than the ones recorded during the baseline period (5.4% respectively). However, it should be noted the sample sizes on total numbers of dolphin sightings were very different between the two periods.
- 3.7.3. Distribution of dolphins engaged in various activities during the present three-month period and baseline period is shown in Figure 6. The one dolphin group engaged in feeding activity was sighted at the southeast corner of Lung Kwu Chau, while the two dolphin groups engaged in socializing activities were both located to the north of Lung Kwu Chau (Figure 6).
- 3.7.4. When compared to the baseline period, distribution of various dolphin activities during the present impact phase monitoring period was drastically different with a much more restricted area of occurrences (Figure 6).
- 3.7.5. Notably, one group of a single dolphin was found to be associated with an operating purse-seiner adjacent to Lung Kwu Chau within the marine park during the present impact phase period.
- *3.8. Summary of photo-identification works*
- 3.8.1. From September to November 2017, over 2,500 digital photographs of Chinese White Dolphins were taken during the impact phase monitoring surveys for the photo-identification work.
- 3.8.2. In total, 23 individuals sighted 42 times altogether were identified (see summary table in Appendix III and photographs of identified individuals in Appendix IV). All of these



re-sightings were made in NWL. Six individuals (i.e. CH34, NL33, NL46, NL49, NL320, NL322, NL328 and WL05) were re-sighted twice, while four other individuals (i.e. NL136, NL182, NL202 and NL286) were re-sighted 3-4 times during the three-month period (Appendix III).

- 3.8.3. Notably, ten of these 23 individuals (i.e. CH34, NL12, NL49, NL104, NL136, NL182, NL202, NL320, NL321 and WL05) were also sighted in Northwest Lantau during the HKBCF monitoring surveys under the same three-month period. Moreover, six individuals (i.e. CH34, NL12, NL49, NL182, NL210 and WL05) were also sighted in West Lantau waters during the HKLR09 monitoring surveys from September to November 2017, showing their extensive individual movements across different survey areas.
- 3.9. Individual range use
- 3.9.1. Ranging patterns of the 23 individuals identified during the three-month study period were determined by fixed kernel method, and are shown in Appendix V.
- 3.9.2. All identified dolphins sighted in the present quarter were utilizing NWL waters only, but have completely avoided NEL waters where many of them have utilized as their core areas in the past (Appendix V). This is in contrary to the extensive movements between NEL and NWL survey areas observed in the earlier impact monitoring quarters as well as the baseline period.
- 3.9.3. On the other hand, three individuals (i.e. NL12, NL182 and NL210) consistently utilized North Lantau waters in the past have extended their range use to WL during the present quarter.
- 3.9.4. In the upcoming quarters, individual range use and movements should be continuously monitored to examine whether there has been any consistent shifts of individual home ranges from North Lantau to West or Southwest Lantau, as such shift could possibly be related to the HZMB-related construction works (see Hung 2017).

#### 4. Conclusion

- 4.1. During this quarter of dolphin monitoring, no adverse impact from the activities of the TMCLKL construction project on Chinese White Dolphins was noticeable from general observations.
- 4.2. Although the dolphins infrequently occurred along the alignment of TMCLKL southern connection viaduct in the past and during the baseline monitoring period, it is apparent that dolphin usage has been significantly reduced in NEL, and many individuals have shifted away from the important habitat around the Brothers Islands.
- 4.3. It is critical to monitor the dolphin usage in North Lantau region in the upcoming quarters, to determine whether the dolphins are continuously affected by the various construction activities in relation to the HZMB-related works, and whether suitable mitigation measure




## 香港鯨豚研究計劃

can be applied to revert the situation.

#### 5. References

- Buckland, S. T., Anderson, D. R., Burnham, K. P., Laake, J. L., Borchers, D. L., and Thomas, L. 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, London.
- Hung, S. K. 2017. Monitoring of marine mammals in Hong Kong waters data collection: final report (2016-17). An unpublished report submitted to the Agriculture, Fisheries and Conservation Department of Hong Kong SAR Government, 162 pp.
- Jefferson, T. A. 2000. Population biology of the Indo-Pacific hump-backed dolphin in Hong Kong waters. Wildlife Monographs 144:1-65.






Figure 1. Distribution of Chinese white dolphin sighting in Northwest and Northeast Lantau during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)




Figure 2. Distribution of Chinese white dolphin sightings in Northwest and Northeast Lantau during the past five autumn quarters (September-November) of HKLR03 impact phase in 2013-17




Figure 3. Distribution of Chinese white dolphins with larger group sizes during HKLR03 impact phase (top) and baseline monitoring surveys (bottom) (green dots: group sizes of 5 or more; purple dots: group sizes of 10 or more)




Figure 4a. Sighting density of Chinese white dolphins with corrected survey effort per km<sup>2</sup> in Northeast and Northwest Lantau survey areas, using data collected during HKLR03 impact monitoring period monitoring period (Sep-Nov 17) (SPSE = no. of on-effort sightings per 100 units of survey effort)




Figure 4b. Density of Chinese white dolphins with corrected survey effort per km<sup>2</sup> in Northeast and Northwest Lantau survey areas, using data collected during HKLR03 impact monitoring period (Sep-Nov 17) (DPSE = no. of dolphins per 100 units of survey effort)




Figure 5. Comparison of density of Chinese white dolphins with corrected survey effort per km<sup>2</sup> in Northwest and Northeast Lantau survey area between the impact monitoring period (September-November 2017) and baseline monitoring period (September-November 2011) (DPSE = no. of dolphins per 100 units of survey effort)




Figure 6. Distribution of Chinese white dolphins engaged in feeding (purple dots), socializing (pink dots) and traveling (green dots) activities during HKLR03 impact phase (top) and baseline monitoring surveys (bottom)

#### Appendix I. HKLR03 Survey Effort Database (September-November 2017)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

DATE	AREA	BEAU	EFFORT	SEASON	VESSEL	TYPE	P/S
15-Sep-17	NW LANTAU	2	26.51	AUTUMN	STANDARD36826	HKLR	Р
15-Sep-17	NW LANTAU	2	10.09	AUTUMN	STANDARD36826	HKLR	S
15-Sep-17	NW LANTAU	3	1.20	AUTUMN	STANDARD36826	HKLR	S
15-Sep-17	NE LANTAU	2	34.49	AUTUMN	STANDARD36826	HKLR	Р
15-Sep-17	NE LANTAU	3	2.20	AUTUMN	STANDARD36826	HKLR	Р
15-Sep-17	NE LANTAU	2	12.01	AUTUMN	STANDARD36826	HKLR	S
18-Sep-17	NW LANTAU	2	28.84	AUTUMN	STANDARD36826	HKLR	Р
18-Sep-17	NW LANTAU	3	7.20	AUTUMN	STANDARD36826	HKLR	Р
18-Sep-17	NW LANTAU	2	12.96	AUTUMN	STANDARD36826	HKLR	S
22-Sep-17	NW LANTAU	1	6.05	AUTUMN	STANDARD36826	HKLR	Р
22-Sep-17	NW LANTAU	2	18.48	AUTUMN	STANDARD36826	HKLR	Р
22-Sep-17	NW LANTAU	3	0.56	AUTUMN	STANDARD36826	HKLR	Р
22-Sep-17	NW LANTAU	1	1.58	AUTUMN	STANDARD36826	HKLR	S
22-Sep-17		2	9.25	AUTUMN	STANDARD36826	HKLR	S
22-Sep-17	NE LANTAU	2	4.68	AUTUMN	STANDARD36826	HKLR	Р
22-Sep-17	NE LANTAU	3	31.06	AUTUMN	STANDARD36826	HKLR	P
22-Sep-17		2	3.30	AUTUMN	STANDARD36826	HKLR	S
22-Sep-17		3	9.06	AUTUMN	STANDARD36826	HKLR	S
29-Sep-17		1	3.40	AUTUMN	STANDARD36826	HKLR	P
29-Sep-17	NW LANTAU	2	13.70	AUTUMN	STANDARD36826	HKLR	Р
29-Sep-17		3	12.90	AUTUMN	STANDARD36826	HKLR	Р
29-Sep-17		4	5.60	AUTUMN	STANDARD36826	HKLR	Р
29-Sep-17	NW LANTAU	2	1.15	AUTUMN	STANDARD36826	HKLR	S
29-Sep-17	NW LANTAU	3	10.06	AUTUMN	STANDARD36826	HKLR	S
4-Oct-17	NW LANTAU	2	0.88	AUTUMN	STANDARD36826	HKLR	P
4-Oct-17	NW LANTAU	3	20.90	AUTUMN	STANDARD36826	HKLR	Р
4-Oct-17	NW LANTAU	4	2.00	AUTUMN	STANDARD36826	HKLR	Р
4-Oct-17	NW LANTAU	2	3.80	AUTUMN	STANDARD36826	HKLR	S
4-Oct-17	NW LANTAU	3	5.02	AUTUMN	STANDARD36826	HKLR	S
4-Oct-17	NW LANTAU	4	2.40	AUTUMN	STANDARD36826	HKLR	S
4-Oct-17	NE LANTAU	2	8.22	AUTUMN	STANDARD36826	HKLR	P
4-Oct-17 4-Oct-17	NE LANTAU	3	11.59	AUTUMN	STANDARD36826	HKLR	P
4-Oct-17	NE LANTAU	2	9.49	AUTUMN	STANDARD36826	HKLR	S
4-Oct-17 4-Oct-17	NE LANTAU	3	1.30	AUTUMN	STANDARD36826	HKLR	S
9-Oct-17	NW LANTAU	2	1.68	AUTUMN	STANDARD36826	HKLR	P
9-Oct-17	NW LANTAU	3	30.32	AUTUMN	STANDARD36826	HKLR	Р
9-Oct-17	NW LANTAU	4	2.50	AUTUMN	STANDARD36826	HKLR	Р
9-Oct-17	NW LANTAU	2	2.30	AUTUMN	STANDARD36826	HKLR	S
9-Oct-17	NW LANTAU	3	4.90	AUTUMN	STANDARD36826	HKLR	S
9-Oct-17	NW LANTAU	4	6.70	AUTUMN	STANDARD36826	HKLR	S
9-Oct-17	NE LANTAU	3	6.99	AUTUMN	STANDARD36826	HKLR	Р
9-Oct-17	NE LANTAU	4	9.93	AUTUMN	STANDARD36826	HKLR	Р
9-Oct-17	NE LANTAU	3	6.79	AUTUMN	STANDARD36826	HKLR	S
9-Oct-17	NE LANTAU	4	3.09	AUTUMN	STANDARD36826	HKLR	S
18-Oct-17	NW LANTAU	2	11.46	AUTUMN	STANDARD36826	HKLR	Р
18-Oct-17	NW LANTAU	3	20.72	AUTUMN	STANDARD36826	HKLR	Р
18-Oct-17	NW LANTAU	2	8.55	AUTUMN	STANDARD36826	HKLR	S
18-Oct-17	NW LANTAU	3	2.50	AUTUMN	STANDARD36826	HKLR	S
18-Oct-17	NE LANTAU	1	2.44	AUTUMN	STANDARD36826	HKLR	Р
18-Oct-17	NE LANTAU	2	27.42	AUTUMN	STANDARD36826	HKLR	Р

### Appendix I. (cont'd)

(Abbreviations: BEAU = Beaufort Sea State; P = Primary Line Effort; S = Secondary Line Effort)

DATE	AREA	BEAU	EFFORT	SEASON	VESSEL	TYPE	P/S
18-Oct-17	NE LANTAU	3	5.50	AUTUMN	STANDARD36826	HKLR	Р
18-Oct-17	NE LANTAU	1	1.70	AUTUMN	STANDARD36826	HKLR	S
18-Oct-17	NE LANTAU	2	11.34	AUTUMN	STANDARD36826	HKLR	S
26-Oct-17	NW LANTAU	2	24.70	AUTUMN	STANDARD36826	HKLR	Р
26-Oct-17	NW LANTAU	3	4.44	AUTUMN	STANDARD36826	HKLR	Р
26-Oct-17	NW LANTAU	2	11.91	AUTUMN	STANDARD36826	HKLR	S
26-Oct-17	NW LANTAU	3	0.85	AUTUMN	STANDARD36826	HKLR	S
1-Nov-17	NW LANTAU	2	17.00	AUTUMN	STANDARD36826	HKLR	Р
1-Nov-17	NW LANTAU	3	15.32	AUTUMN	STANDARD36826	HKLR	Р
1-Nov-17	NW LANTAU	2	8.38	AUTUMN	STANDARD36826	HKLR	S
1-Nov-17	NW LANTAU	3	2.53	AUTUMN	STANDARD36826	HKLR	S
1-Nov-17	NE LANTAU	2	29.72	AUTUMN	STANDARD36826	HKLR	Р
1-Nov-17	NE LANTAU	3	5.10	AUTUMN	STANDARD36826	HKLR	Р
1-Nov-17	NE LANTAU	2	10.07	AUTUMN	STANDARD36826	HKLR	S
1-Nov-17	NE LANTAU	3	2.41	AUTUMN	STANDARD36826	HKLR	S
8-Nov-17	NW LANTAU	2	13.77	AUTUMN	STANDARD36826	HKLR	Р
8-Nov-17	NW LANTAU	3	14.05	AUTUMN	STANDARD36826	HKLR	Р
8-Nov-17	NW LANTAU	2	10.58	AUTUMN	STANDARD36826	HKLR	S
8-Nov-17	NW LANTAU	3	1.80	AUTUMN	STANDARD36826	HKLR	S
17-Nov-17	NW LANTAU	2	8.53	AUTUMN	STANDARD36826	HKLR	Р
17-Nov-17	NW LANTAU	3	18.98	AUTUMN	STANDARD36826	HKLR	Р
17-Nov-17	NW LANTAU	2	9.37	AUTUMN	STANDARD36826	HKLR	S
17-Nov-17	NW LANTAU	3	3.55	AUTUMN	STANDARD36826	HKLR	S
24-Nov-17	NW LANTAU	2	3.81	AUTUMN	STANDARD36826	HKLR	Р
24-Nov-17	<b>NW LANTAU</b>	3	28.72	AUTUMN	STANDARD36826	HKLR	Р
24-Nov-17	<b>NW LANTAU</b>	2	4.40	AUTUMN	STANDARD36826	HKLR	S
24-Nov-17	<b>NW LANTAU</b>	3	6.27	AUTUMN	STANDARD36826	HKLR	S
24-Nov-17	NE LANTAU	2	30.83	AUTUMN	STANDARD36826	HKLR	Р
24-Nov-17	NE LANTAU	3	4.97	AUTUMN	STANDARD36826	HKLR	Р
24-Nov-17	NE LANTAU	1	1.20	AUTUMN	STANDARD36826	HKLR	S
24-Nov-17	NE LANTAU	2	10.10	AUTUMN	STANDARD36826	HKLR	S

Appendix II. HKLR03 Chinese White Dolphin Sighting Database (September-November 2017)


(Abberviations: STG# = Sighting Number; HRD SZ = Dolphin Herd Size; BEAU = Beaufort Sea State; PSD = Perpendicular Distance; BOAT ASSOC. = Fishing Boat Association P/S: Sighting Made on Primary/Secondary Lines)

DATE	STG#	TIME	HRD SZ	AREA	BEAU	PSD	EFFORT	TYPE	NORTHING	EASTING	SEASON	BOAT ASSOC.	P/S
22-Sep-17	1	1152	6	NW LANTAU	2	320	ON	HKLR	823991	807501	AUTUMN	NONE	Р
22-Sep-17	2	1244	3	NW LANTAU	1	250	ON	HKLR	825349	809502	AUTUMN	NONE	Р
29-Sep-17	1	1309	2	NW LANTAU	4	140	ON	HKLR	827215	806416	AUTUMN	NONE	Р
4-Oct-17	1	1143	5	NW LANTAU	3	52	ON	HKLR	828985	807490	AUTUMN	NONE	Р
18-Oct-17	1	1149	1	NW LANTAU	2	65	ON	HKLR	826905	806487	AUTUMN	NONE	Р
18-Oct-17	2	1159	1	NW LANTAU	2	264	ON	HKLR	825632	806485	AUTUMN	PURSE-SEINE	Р
26-Oct-17	1	1135	1	NW LANTAU	2	34	ON	HKLR	826737	807455	AUTUMN	NONE	Р
1-Nov-17	1	1126	6	NW LANTAU	3	371	ON	HKLR	830641	804652	AUTUMN	NONE	Р
1-Nov-17	2	1152	8	NW LANTAU	2	529	ON	HKLR	827437	806499	AUTUMN	NONE	Р
8-Nov-17	1	1129	2	NW LANTAU	2	317	ON	HKLR	826272	807434	AUTUMN	NONE	Р
17-Nov-17	1	1155	12	NW LANTAU	2	627	ON	HKLR	829665	805381	AUTUMN	NONE	S
24-Nov-17	1	1023	2	NW LANTAU	3	8	ON	HKLR	816588	804674	AUTUMN	NONE	Р
24-Nov-17	2	1155	1	NW LANTAU	3	0	ON	HKLR	826850	806436	AUTUMN	NONE	Р


## Appendix III. Individual dolphins identified during HKLR03 monitoring surveys in September - November 2017

ID#	DATE	STG#	AREA
CH34	01/11/17	2	NW LANTAU
	17/11/17	1	NW LANTAU
NL12	04/10/17	1	NW LANTAU
NL33	01/11/17	2	NW LANTAU
	17/11/17	1	NW LANTAU
NL46	22/09/17	1	NW LANTAU
	17/11/17	1	NW LANTAU
NL49	22/09/17	1	NW LANTAU
	17/11/17	1	NW LANTAU
NL104	04/10/17	1	NW LANTAU
NL123	22/09/17	1	NW LANTAU
NL136	04/10/17	1	NW LANTAU
	18/10/17	2	NW LANTAU
	01/11/17	2	NW LANTAU
	08/11/17	1	NW LANTAU
NL145	17/11/17	1	NW LANTAU
NL182	04/10/17	1	NW LANTAU
	18/10/17	1	NW LANTAU
	01/11/17	2	NW LANTAU
	24/11/17	2	NW LANTAU
NL202	22/09/17	2	NW LANTAU
	29/09/17	1	NW LANTAU
	01/11/17	2	NW LANTAU
NL210	01/11/17	2	NW LANTAU
NL242	22/09/17	1	NW LANTAU
NL261	17/11/17	1	NW LANTAU
NL272	17/11/17	1	NW LANTAU
NL286	22/09/17	2	NW LANTAU
	29/09/17	1	NW LANTAU
	01/11/17	2	NW LANTAU
	17/11/17	1	NW LANTAU
NL296	22/09/17	1	NW LANTAU
NL320	01/11/17	2	NW LANTAU
	17/11/17	1	NW LANTAU
NL321	04/10/17	1	NW LANTAU
NL322	01/11/17	2	NW LANTAU
	17/11/17	1	NW LANTAU
NL328	08/11/17	1	NW LANTAU
	17/11/17	1	NW LANTAU
WL05	22/09/17	1	NW LANTAU
	17/11/17	1	NW LANTAU
WL145	24/11/17	1	NW LANTAU


Appendix IV. Twenty-three individual dolphins that were identified during September to November 2017 under HKLR03 impact phase monitoring surveys




Appendix IV. (cont'd)



Appendix IV. (cont'd)




Appendix IV. (cont'd)




Appendix IV. (cont'd)




Appendix IV. (cont'd)




Appendix V. Ranging patterns (95% kernel ranges) of 23 individual dolphins that were sighted during HKLR03 impact phase monitoring period (note: yellow dots indicate sightings made in September – November 2017 during HKLR03 and HKLR09 monitoring surveys; the yellow dots with the red circles indicate the ones made during HKBCF monitoring surveys)



Appendix V. (cont'd)



Appendix V. (cont'd)



Appendix V. (cont'd)

